

## Analytic Solutions

1. For the IVP

$$(1-x)y'' + xy' - y = 0, \quad y(0) = -3, y'(0) = 2$$

(a) Determine the minimum radius of convergence of solutions around  $x_0 = 0$ .

$$P = 1-x \Rightarrow P(1) = 0$$

$$\Downarrow \\ R_{oC} = (1) - (0) = \boxed{1}$$

(b) If  $y = \phi(x)$  is a solution of the IVP, find  $\phi''(0)$ ,  $\phi'''(0)$ , and  $\phi^{(4)}(0)$ .

$$\phi(0) = -3$$

$$\phi'(0) = 2$$

$$(1-x)y''' - 1y'' + xy' + 1y = 0$$

$$\phi'''(0) = 1 \cdot \phi''(0) = \boxed{-3}$$

$$(1-x)y'' + xy' - y = 0$$

$$y'' = \frac{-x}{1-x} y' + \frac{1}{1-x} y$$

$$\phi''(0) = 0\phi'(0) + 1\phi(0) \\ = \boxed{-3}$$

$$(1-x)y^{(4)} + (-1-1+x)y''' + 1y'' = 0$$

$$\phi^{(4)}(0) = 2\phi'''(0) - 1\phi''(0) \\ = -6 + 3 = \boxed{-3}$$

(c) Write down the first five terms of the analytic power series solution

$$y = \sum_{n=0}^{\infty} a_n x^n$$

by using the relationship  $n!a_n = \phi^{(n)}(x_0)$ .

$$a_0 = y(0) = -3$$

$$a_1 = y'(0) = 2$$

$$a_2 = \frac{1}{2} y''(0) = -\frac{3}{2}$$

$$a_3 = \frac{1}{6} y'''(0) = -\frac{1}{2}$$

$$a_4 = \frac{1}{24} y^{(4)}(0) = -\frac{1}{8}$$

$$y(x) = -3 + 2x - \frac{3}{2}x^2 - \frac{1}{2}x^3 - \frac{1}{8}x^4 + \dots$$

## More Power Series

2. Find a general solution to the following differential equation using the power series method.

$$y'' + xy = 0, \quad y(0) = 1, y'(0) = 0.$$

$$\downarrow \\ x_0 = 0$$

$$y = \sum_{n=0}^{\infty} a_n x^n$$

$$y'' = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}$$

$$y'' + xy = 0$$

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} + x \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + \sum_{n=1}^{\infty} a_{n-1} x^n = 0$$

$$(2)(1) a_2 x^0 + \sum_{n=1}^{\infty} ((n+2)(n+1) a_{n+2} + a_{n-1}) x^n = 0$$

↓

$$a_2 = 0$$

$$(n+2)(n+1) a_{n+2} + a_{n-1} = 0 \quad \text{for } n \geq 1$$

$$(n+3)(n+2) a_{n+3} + a_n = 0 \quad \text{for } n \geq 0$$

$$a_{n+3} = -\frac{a_n}{(n+2)(n+3)} \quad \text{for } n \geq 0$$

$$a_0 = y(0) = 1$$

$$a_1 = y'(0) = 0$$

$$a_2 = 0$$

$$\Rightarrow y = 1 - \frac{1}{6} x^3 + \frac{1}{180} x^6 - \dots + \frac{(-1)^n}{2 \cdot 3 \cdot 5 \cdot 6 \cdot \dots \cdot 3n} x^{3n} + \dots$$

$$a_3 = -\frac{1}{2 \cdot 3} = -\frac{1}{6}$$

$$a_6 = +\frac{1/6}{5 \cdot 6} = \frac{1}{180}$$

$$a_{3n} = \frac{(-1)^n}{2 \cdot 3 \cdot 5 \cdot 6 \cdot \dots \cdot (3n-1)(3n)}$$

## Euler Equations

3. Solve the Euler equation  $2x^2y'' + 3xy' - y = 0$ ,  $x > 0$  by looking for solutions of the form  $y = x^r$ .

$$y = x^r$$

$$y' = r x^{r-1}$$

$$y'' = r(r-1)x^{r-2}$$

$$2x^2 r(r-1)x^{r-2} + 3x r x^{r-1} - x^r = 0$$

$$2r(r-1)x^r + 3r x^r - x^r = 0$$

$$(2r(r-1) + 3r - 1) \cancel{x^r}^{x>0} = 0$$

$$2r^2 - 2r + 3r - 1 = 0$$

$$2r^2 + r - 1 = 0$$

$$(2r-1)(r+1) = 0$$

$$r = \frac{1}{2}, -1$$

↓

$$y = C_1 \sqrt{x} + \frac{C_2}{x}$$

- (a) Use the Wronskian to show that the two solutions are linearly independent for  $x > 0$ .

$$W\left(\sqrt{x}, \frac{1}{x}\right) = \begin{vmatrix} \sqrt{x} & \frac{1}{x} \\ \frac{1}{2\sqrt{x}} & -\frac{1}{x^2} \end{vmatrix} = -x^{-3/2} - \frac{1}{2}x^{-3/2}$$

$$= -\frac{3}{2}x^{-3/2} \neq 0 \quad \forall x > 0$$

↓

l. i.

4. For the Euler equation,

$$x^2 y'' + 5xy' + 4y = 0, \quad x > 0,$$

(a) Find one solution  $y_1(x)$  by making the substitution  $y = x^r$ .

$$\begin{array}{l} y = x^r \\ y' = r x^{r-1} \\ y'' = r(r-1)x^{r-2} \end{array} \quad \left| \quad \begin{array}{l} x^2 r(r-1)x^{r-2} + 5x r x^{r-1} + 4x^r = 0 \\ r(r-1) + 5r + 4 = 0 \\ r = -2 \quad \Rightarrow \quad y_1 = \frac{1}{x^2} \end{array} \right.$$

(b) Use the method of reduction of order to find the other solution:

i. Assume a second solution of the form  $y_2(x) = u(x)y_1(x)$ . Plug into the differential equation and simplify to an equation involving  $u''$  and  $u'$ .

ii. Solve for  $u'$ ,

iii. Antidifferentiate to determine  $u$ .

$$\begin{array}{l} y_2 = u y_1 \\ y_2' = u' y_1 + u y_1' \\ y_2'' = u'' y_1 + 2u' y_1' + u y_1'' \end{array} \quad \left| \quad \begin{array}{l} x^2 (u'' y_1 + 2u' y_1' + u y_1'') + 5x (u' y_1 + u y_1') + 4u y_1 = 0 \\ \underbrace{x^2 y_1}_1 u'' + \underbrace{(2x^2 y_1' + 5x y_1)}_{-\frac{4}{x} + \frac{5}{x}} u' + \underbrace{(x^2 y_1'' + 5x y_1' + 4y_1)}_0 u = 0 \\ u'' + \frac{1}{x} u' = 0 \end{array} \right.$$

$$(x u')' = 0 \cdot x = 0$$

$$x u' = c$$

$$u = \int \frac{1}{x} dx = \ln x + c$$

$$y_2 = (\ln x) \cdot \frac{1}{x^2}$$

$$\boxed{y = \frac{c_1}{x^2} + \frac{c_2 \ln x}{x^2}}$$